



Outcomes from sudden cardiac arrest in US high schools: a 2-year prospective study from the National Registry for AED Use in Sports

Jonathan A Drezner, Brett G Toresdahl, Ashwin L Rao, et al.

Br J Sports Med published online October 11, 2013
doi: 10.1136/bjsports-2013-092786

Updated information and services can be found at:
<http://bjsm.bmj.com/content/early/2013/10/11/bjsports-2013-092786.full.html>

These include:

- | | |
|-------------------------------|---|
| References | This article cites 34 articles, 15 of which can be accessed free at: http://bjsm.bmj.com/content/early/2013/10/11/bjsports-2013-092786.full.html#ref-list-1 |
| P<P | Published online October 11, 2013 in advance of the print journal. |
| Email alerting service | Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article. |

Notes

Advance online articles have been peer reviewed, accepted for publication, edited and typeset, but have not yet appeared in the paper journal. Advance online articles are citable and establish publication priority; they are indexed by PubMed from initial publication. Citations to Advance online articles must include the digital object identifier (DOIs) and date of initial publication.

To request permissions go to:
<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:
<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:
<http://group.bmj.com/subscribe/>

Outcomes from sudden cardiac arrest in US high schools: a 2-year prospective study from the National Registry for AED Use in Sports

Jonathan A Drezner,¹ Brett G Toresdahl,¹ Ashwin L Rao,¹ Ella Huszti,² Kimberly G Harmon¹

¹Department of Family Medicine, University of Washington, Seattle, Washington, USA

²Harborview Center for Prehospital Emergency Care, University of Washington, Seattle, Washington, USA

Correspondence to

Dr Jonathan A Drezner, Department of Family Medicine, University of Washington, P.O. Box 356040, Seattle, WA 98195, USA; jdrezner@uw.edu

Accepted 24 September 2013

ABSTRACT

Background Sudden cardiac arrest (SCA) is the leading cause of death in athletes during exercise. The effectiveness of school-based automated external defibrillator (AED) programmes has not been established through a prospective study.

Methods A total of 2149 high schools participated in a prospective observational study beginning 1 August 2009, through 31 July 2011. Schools were contacted quarterly and reported all cases of SCA. Of these 95% of schools confirmed their participation for the entire 2-year study period. Cases of SCA were reviewed to confirm the details of the resuscitation. The primary outcome was survival to hospital discharge.

Results School-based AED programmes were present in 87% of participating schools and in all but one of the schools reporting a case of SCA. Fifty nine cases of SCA were confirmed during the study period including 26 (44%) cases in students and 33 (56%) in adults; 39 (66%) cases occurred at an athletic facility during training or competition; 55 (93%) cases were witnessed and 54 (92%) received prompt cardiopulmonary resuscitation. A defibrillator was applied in 50 (85%) cases and a shock delivered onsite in 39 (66%). Overall, 42 of 59 (71%) SCA victims survived to hospital discharge, including 22 of 26 (85%) students and 20 of 33 (61%) adults. Of 18 student-athletes 16 (89%) and 8 of 9 (89%) adults who arrested during physical activity survived to hospital discharge.

Conclusions High school AED programmes demonstrate a high survival rate for students and adults who suffer SCA on school campus. School-based AED programmes are strongly encouraged.

INTRODUCTION

Sudden cardiac arrest (SCA) is the leading cause of death in the USA and afflicts over 350 000 persons annually.¹ Survival following out-of-hospital cardiac arrest is critically dependent on prompt recognition, early cardiopulmonary resuscitation (CPR) and access to early defibrillation.²⁻³ Public access defibrillation (PAD) programmes are designed to shorten the time interval from cardiac arrest to defibrillation by the strategic placement of automated external defibrillators (AEDs) in public locations. Several studies have demonstrated improved survival with use of AED by trained or untrained lay responders within the first minutes following SCA.⁴⁻⁹

SCA is also the leading cause of sudden death in young athletes during exercise.¹⁰⁻¹¹ The risk of SCA is increased during physical activity in

individuals of all ages with underlying cardiovascular disorders.¹²⁻¹⁵ The tragic and unexpected death of a young person during sport remains a highly visible and devastating event for the general public and the medical community and invariably stimulates review of emergency preparations at athletic events. The placement of AEDs in schools and at athletic venues has become the cornerstone of emergency response planning and the prevention of sudden cardiac death in young athletes.¹⁶⁻¹⁸ School-based AED programmes provide a means of early defibrillation for students as well as adults who suffer SCA on school campus. A study of 1710 high schools equipped with onsite AEDs reported a 64% survival rate in student-athletes as well as adults who suffered SCA on school campus.¹⁷ However, limitations of the study were its cross-sectional design and potential for responder bias to influence the results.

The purpose of this study was to prospectively monitor a large cohort of US high schools to determine the outcomes of SCA in schools and the potential effectiveness of school-based AED programmes.

METHODS

Study design

High schools enrolled in the National Registry for AED Use in Sports participated in a prospective observational study from 1 August 2009 to 31 July 2011. The National Registry for AED Use in Sports was developed in 2005 to monitor emergency planning and outcomes from SCA in the university athletic setting and expanded in 2006 to monitor SCA in US high schools. The registry consists of a web-based questionnaire and database management system including a comprehensive survey on emergency response planning for SCA and additional questions regarding the details of any SCA or AED use.

At the beginning of the study, each school confirmed their participation and was asked to update their survey on emergency response planning. Schools were asked to report any case of SCA or AED use that occurred on school campus. Schools were contacted quarterly by email as a reminder throughout the study period. At the conclusion of the study, each school was asked to confirm their participation in the 2-year study and reaffirm the number of cases of SCA or AED use (if any) reported. Non-responding schools were contacted by email on three additional occasions. For all non-responding schools, attempts were made to contact

To cite: Drezner JA, Toresdahl BG, Rao AL, *et al.* *Br J Sports Med* Published Online First: [please include Day Month Year] doi:10.1136/bjsports-2013-092786

each school directly by phone to confirm their participation in the 2-year study and document any cases of SCA or AED use.

For every reported case of SCA, the details of the case were reviewed directly with the school to ensure the accuracy of the information provided and that the case occurred during the study period. Details of the resuscitation were reviewed with a school staff member familiar with the emergency response, such as a school administrator, athletic director, nurse, coach or certified athletic trainer. In some instances, publicly available media reports of the incident were reviewed to supplement the information provided. Potential cases of SCA were included only if the event occurred on school campus and within the study period, was non-traumatic (except commotio cordis) and of primary cardiac origin and if the victim was confirmed unconscious with absent pulse and respirations.

Study outcomes

The primary outcome measure was survival to hospital discharge. Additional data variables collected included witnessed arrest, provision of bystander CPR, school AED use, responding emergency medical services (EMS) defibrillator use, shock deployment, location of SCA and the age, sex, ethnicity and sport (if applicable) of the victim.

Statistical analysis

Descriptive statistics such as proportions and means were used to describe the data, as appropriate. A post hoc univariate logistic regression analysis was performed to determine factors associated with survival to hospital discharge. Specifically, we compared survival rates in schools with onsite AED use versus schools where the defibrillator used was brought by responding EMS. We also compared survival in schools with an established emergency action plan for SCA versus schools with no emergency action plan. The study was not powered to compare schools with and without an AED programme. This study was approved by the Human Subjects Division at the University of Washington.

RESULTS

Participants

A total of 2149 high schools distributed throughout all 50 states participated in the study. The mean school enrolment was 963 and the mean number of student-athletes per school was 367. Schools were 86% public and 14% private and the location of the schools was 48% rural, 33% suburban, 15% urban and 4% inner city (table 1).

Of the 2149 schools 2045 (95%) confirmed their participation for the entire 2-year study period. The remaining 104 (5%) school representatives could not be reached after multiple attempts by email and phone.

Emergency planning

School-based AED programmes were present in 87% of participating schools. Of the schools with at least one AED, the mean number of devices per school was 3.1. Eighty-three per cent of schools had a written emergency action plan for SCA. Of schools with an emergency action plan, 67% consulted the local EMS when developing the plan. School personnel most likely to be trained in CPR were coaches (80% of schools), school nurses (77%), certified athletic trainers (68%) and administrators (65%). School personnel most likely to be trained in AED use were coaches (75% of schools), school nurses (74%), certified athletic trainers (65%) and administrators (65%). Only 45% of schools reviewed and practiced their emergency response to

Table 1 School demographics

| School demographics (%) | |
|-------------------------|----|
| Setting | |
| Inner city | 4 |
| Urban | 15 |
| Suburban | 33 |
| Rural | 48 |
| Type | |
| Public | 86 |
| Private | 14 |
| Student enrolment | |
| Less than 500 | 37 |
| 501 to 1000 | 26 |
| 1001 to 1500 | 15 |
| 1501 to 2000 | 11 |
| Greater than 2000 | 11 |
| Student-athletes | |
| Less than 200 | 36 |
| 201 to 400 | 29 |
| 401 to 600 | 17 |
| Greater than 600 | 18 |

SCA at least once annually and only 20% of schools posted their emergency action plan at athletic venues.

Of the schools without AEDs, 263 schools indicated one or more obstacles to obtaining AEDs. The most common reasons for not having AEDs were limited financial resources (80%), medicolegal concerns (24%), uncertainty on where to place AEDs (13%), school policy (10%) and unproven cost-effectiveness (1%).

SCA cases

A total of 129 possible cases of SCA were reported. Of these, 59 cases of SCA were confirmed and included for further analysis; 69 cases were excluded: of these 32 occurred outside the study time period, 20 were determined not to be SCA, 13 occurred outside the school campus and 4 were induced by trauma (eg, head injury). One case could not be confirmed as the school representative and first responders could not be contacted. The annual incidence of SCA occurring in a high school was 1 in 73 schools/year.

Case demographics

Twenty-six (44%) cases occurred in students and 33 (56%) occurred in adults. The mean age was 16 years (range 13–20) in students and 56 years (range 20–80) in adults. Of the cases occurring in students, 18 (69%) were student-athletes and 8 (31%) non-athletes. Three (17%) of the cases in student-athletes were from commotio cordis (1 each in football, ice hockey and lacrosse). Of the cases occurring in adults, 10 (30%) were spectators, 9 (27%) visitors on campus, 7 (21%) school staff, 4 (12%) athletic officials and 3 (9%) coaches.

Forty-seven (80%) cases occurred in males. Of the cases occurring in student-athletes, 16 (89%) were male and 2 (11%) were female. SCA occurred equally in males and females in the eight cases of student non-athletes. Of the cases occurring in adults, 27 (82%) were male and 6 (18%) were female.

The ethnicity of persons suffering SCA was 49 (82%) Caucasian, 8 (14%) African-American/black, and 2 (4%) Hispanic. The ethnicity of student-athletes with SCA was 12

(67%) Caucasian, 5 (28%) African-American/black and 1 (6%) Hispanic.

Location and setting

Athletic facilities were the most common location of SCA, comprising 66% of cases (box 1). Twenty (34%) cases were during games and 19 (32%) cases were during practice or training. Twenty-seven (46%) cases were in persons participating in physical activity on the school campus such as student-athletes, officials and visitors using athletic facilities. Basketball and football were the most frequent sporting events in which SCA occurred with 13 and 9 cases, respectively. Other sports in which SCA occurred included baseball (3), track and field (3), ice hockey (2), soccer (2), softball (2), cheerleading (1), cross country (1), lacrosse (1), swimming (1), and wrestling (1). Of the cases that did not occur at a sporting event, 16 (27%) occurred in a classroom building and 4 (7%) occurred in a parking lot.

Emergency response

SCA was witnessed in 55 (93%) of overall cases and in all cases involving a student-athlete. The presence or absence of seizure-like activity was reported in 40 cases, with 36% of adults and 38% of student-athletes reported to have brief seizure-like activity after collapse. The most frequent school responders to the

event were administrative staff (22), certified athletic trainers (21), lay visitors (19), school nurses (19), coaches (17) and teachers (17). EMS personnel were stationed onsite at the athletic venue prior to the SCA in three cases. The average estimated time to arrival of EMS when not pre-stationed onsite was 7.8 min. CPR was provided by lay responders or onsite EMS personnel in 54 (92%) cases.

School-based AED programmes were present in all but one of the schools reporting a case of SCA. A defibrillator was brought to the resuscitation in all cases. The source of the defibrillator used in the resuscitation was located onsite at the school in 41 (70%) cases, brought by responding offsite EMS in 15 (25%) cases and in 3 (5%) cases EMS was already onsite and provided the defibrillator. Application of the defibrillator was confirmed in 50 (85%) cases and a shock delivered onsite in 39 (66%) cases.

Survival

Overall, 42 of 59 (71%) SCA victims survived to hospital discharge, including 20 of 33 (61%) adults and 22 of 26 (85%) students (table 2). Survival in males was 75% and 58% in females. Of 18 student-athletes 16 (89%) survived to hospital discharge, as did 8 of 9 (89%) adults who arrested during physical activity (figure 1). Of the 39 cases where a shock was delivered onsite, 34 (87%) survived.

Eighty per cent of SCA victims survived to hospital discharge if the school supplied the AED used in the resuscitation versus 50% if the defibrillator was brought by responding offsite EMS (unadjusted OR 4.0, 95% CI (1.14 to 14.02), p value 0.03). The survival rate was 79% in schools with an established emergency action plan for SCA versus 44% in those without (unadjusted OR 4.6, 95% CI (1.04 to 20.48), p value 0.03). None of the unwitnessed cases of SCA survived to hospital discharge.

DISCUSSION

The single greatest factor affecting survival from SCA is the time interval from cardiac arrest to defibrillation.¹⁹ In the USA, historical survival rates from out-of-hospital cardiac arrest using conventional EMS systems are less than 8%.^{20–22} Survival following SCA has been greatly improved by lay rescuers and PAD programmes designed to shorten the time interval from SCA to shock delivery.^{4–9} These programmes train lay rescuers and non-traditional first responders in CPR and AED use and place AEDs in public locations where risk for SCA is high.

The PAD trial found survival from out-of-hospital SCA doubled when lay responders were trained and equipped with AEDs compared with CPR alone.⁵ The Resuscitation Outcomes Consortium Epistery Cardiac Arrest registry also demonstrated that bystander application of AED prior to EMS arrival nearly doubled the likelihood of survival following out-of-hospital SCA.⁸ Other studies of rapid defibrillation in specific public settings such as casinos, airlines and airports have shown survival rates ranging from 41% to 74% if bystander CPR was provided and defibrillation occurred within 3–5 min of collapse.^{4 6 7} Essential elements to the success of these programmes include training of anticipated responders in CPR and AED use, a structured and practiced response, and short response times.

This is the first large prospective study of PAD in the school setting. The results of this study demonstrate a high survival rate for victims of SCA on school campus and provide compelling support of school-based AED programmes. The high survival rates in this study are likely related to several factors known to increase the probability of survival from SCA. Over

Box 1 Location of sudden cardiac arrest

Location of sudden cardiac arrest (number of cases)

Athletic facility (39)

Baseball field (3)

Student-athlete (2), spectator (1)

Football stadium (5)

Student-athlete (3), visiting adult athlete (1), spectator (1)

Gymnasium/basketball (13)

Spectator (5), student-athlete (3), visiting adult athlete (3), coach (1), official (1)

Gymnasium/cheerleading (1)

Student-athlete (1)

Gymnasium/other training (3)

Student-athlete (3)

Gymnasium/wrestling (1)

Spectator (1)

Cross country (1)

Student-athlete (1)

Hockey rink (2)

Student-athlete (1), visitor (1)

Lacrosse field (1)

Student athlete (1)

Soccer field (3)

Official (2), coach (1)

Softball field (2)

Spectator (1), official (1)

Swimming pool (1)

Student-athlete (1)

Track and field (3)

Student-athlete (2), coach (1)

Classroom building (16)

Student (8), school staff (5), visitor (3)

Parking lot (4)

School staff (2), spectator (1), visitor (1)


Table 2 Survival from sudden cardiac arrest in US high schools

| | Number of cases | Survival to hospital discharge |
|-------------------------------|-----------------|--------------------------------|
| All cases | 59 | 42 (71%) |
| Gender | | |
| Male | 47 | 35 (75%) |
| Female | 12 | 7 (58%) |
| Ethnicity | | |
| African-American | 8 | 4 (50%) |
| Caucasian | 47 | 34 (72%) |
| Hispanic | 2 | 2 (100%) |
| Other | 2 | 2 (100%) |
| Students | | |
| All students | 26 | 22 (85%) |
| Student-athletes | 18 | 16 (89%) |
| Males | 16 | 14 (88%) |
| Females | 2 | 2 (100%) |
| Student non-athletes | 8 | 6 (75%) |
| Setting | | |
| Sporting event | 40 | 32 (80%) |
| Game/competition | 21 | 16 (76%) |
| Practice/training | 19 | 16 (84%) |
| Non-sporting event | 19 | 10 (53%) |
| Physical activity | | |
| During physical activity | 27 | 24 (89%) |
| Student-athletes | 18 | 16 (89%) |
| Adults | 9 | 8 (89%) |
| Not during physical activity | 32 | 18 (56%) |
| Witnessed | | |
| Witnessed | 55 | 42 (76%) |
| Unwitnessed | 4 | 0 (0%) |
| CPR | | |
| Performed | 54 | 39 (72%) |
| Not performed | 5 | 3 (60%) |
| Source of AED used | | |
| School | 41 | 32 (78%) |
| EMS (offsite) | 16 | 8 (50%) |
| EMS (onsite) | 2 | 2 (100%) |
| Defibrillation onsite | | |
| Shock delivered | 39 | 34 (87%) |
| Shock not indicated | 15 | 6 (40%) |
| AED not applied | 3 | 2 (67%) |
| Unable to confirm | 2 | 0 (0%) |
| Emergency action plan for SCA | | |
| Established | 47 | 37 (79%) |
| Not established | 9 | 4 (44%) |
| Unable to confirm | 3 | 1 (33%) |

AED, automated external defibrillator; CPR, cardiopulmonary resuscitation; EMS, emergency medical services; SCA, sudden cardiac arrest.

90% of cases were witnessed and received prompt CPR, 83% of schools had an established emergency action plan for SCA, school AEDs were readily accessible and used in nearly 70% of cases, and two-thirds of victims received defibrillation onsite.

Cases in which the school had AED but it was not used in the resuscitation (ie, responding EMS provided the defibrillator) also had a reasonably high survival rate. This may reflect that school-based AED programmes predict other important determinants of survival and better overall emergency planning for SCA. In a review of emergency planning for SCA in over 3300

**Figure 1** Survival to hospital discharge in students and adults with sudden cardiac arrest on a high school campus.

high schools, schools with AED programmes were compared to schools without AED programmes and found to be more likely to ensure access to early defibrillation, establish an emergency action plan for SCA, review the emergency action plan at least annually, consult EMS in the development of the emergency action plan, and establish a communication system to activate emergency responders.²³

On any given school day, as much as 20% of the combined US adult and child population can be found in schools.²⁴ Thus, schools have been identified as an important location to consider AED programmes.²⁴ Awareness regarding SCA in children and young athletes also has received increasing attention as a significant cause of mortality.^{11 25 26} A working group of the National Heart, Lung and Blood Institute recently described sudden cardiac death in the young as a critical public health issue.²⁷ The cause of SCA in children and young athletes is typically the result of undiagnosed structural or electrical cardiovascular diseases, such as hypertrophic cardiomyopathy, anomalous coronary arteries and ion channel disorders.^{11 28} Prior studies have reported that survival following exercise-related SCA in the young is poor.^{29 30} This study suggests SCA in students and student-athletes is largely a survivable event through prompt treatment and access to AED.

While there is a growing trend towards AED placement in schools, school-based AED programmes are not yet a universal standard.³¹ The primary obstacle for a school to implement an AED programme is financial concerns. Greater resource allocation towards emergency planning for SCA is needed to ensure widespread access to AEDs in the school setting. Given two-thirds of SCA on a school campus occurs at an athletic venue, prioritising AED placement to these locations should be considered if resources are limited.

This study is limited by the lack of a large control group of SCA cases in schools without an AED programme. With the increasing prevalence of AEDs in schools, establishing a sizeable cohort without AEDs is difficult. However, it is unlikely that the survival rate in schools without an AED programme would be similar. In a retrospective study of EMS treated SCA in schools, the survival rate was 39% in a community with a robust emergency response system.³² Our study also did not follow SCA victims beyond hospital discharge and thus neurological outcomes are unknown. In previous PAD studies that assessed neurological outcome, the functional status of survivors was generally normal.^{4 5} Lastly, although SCA is a highly visible event on a school campus, it is possible some cases of SCA were

not reported. The 95% confirmed 2-year follow-up in this study cohort suggests the potential for missed cases is small.

Past studies have suggested a survival benefit for school-based AED programmes but were limited by their retrospective or cross-sectional designs.^{17 31 33 34} The prospective observational design in this study is similar to prior reports that established the benefit of early defibrillation and subsequently changed safety standards in specific public locations such as casinos, airlines and airports.^{4 6 7} The number of SCA cases in these studies ranged from 21 to 148 and a high survival rate (61–87%) was consistently achieved across different study settings if the victim was found with a shockable rhythm and received defibrillation.^{4 6 7} This study also demonstrated an 87% survival rate to hospital discharge if a shock was deployed.

CONCLUSIONS

High school AED programmes demonstrate a high survival rate for students as well as adults who suffer SCA on school campus. SCA is the leading cause of death in the USA and the leading cause of death in exercising young athletes. School-based AED programmes represent an important public safety measure and an effective strategy for the prevention of sudden cardiac death during sports. Schools are a strategic location to serve large concentrations of people at risk for SCA and school-based AED programmes should be strongly supported.

What are the new findings?

- ▶ Approximately 1 in 70 high schools will have a sudden cardiac arrest (SCA) on campus each year, and nearly half of these events will be in students or student-athletes.
- ▶ SCA in student-athletes is largely a survivable event (>85%) if the event is witnessed, the school has an established emergency action plan and the victim receives prompt cardiopulmonary resuscitation and early defibrillation from an automated external defibrillator (AED).
- ▶ Survival rates are higher in schools with an established emergency action plan for SCA versus those without (79% vs 44%; OR 4.6) and if an onsite AED is used versus an offsite AED provided by emergency medical services (80% vs 50%; OR 4.0).
- ▶ School-based AED programmes represent an important public safety measure and an effective strategy for the prevention of sudden cardiac death during sports.

How might it impact on clinical practice in the near future?

- ▶ Coaches, physical education teachers and other school staff responsible for supervising sports and physical activity in students should be trained in prompt recognition of sudden cardiac arrest (SCA), cardiopulmonary resuscitation and use of an AED.
- ▶ School sponsored sporting events should have an established emergency action plan for SCA and have a rapid access to an AED.
- ▶ School-based public access defibrillation programmes positively impact public safety standards for SCA and should be strongly encouraged.

Contributors JD, BGT, ALR and KGH all contributed to the experimental design, data collection and analysis and preparation of the manuscript. EH contributed to the data and statistical analysis and preparation of the manuscript. All authors have approved the final manuscript.

Funding National Operating Committee on Standards for Athletic Equipment (NOCSAE).

Competing interests None.

Ethics approval University of Washington Human Subjects Division.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- 1 Roger VL, Go AS, Lloyd-Jones DM, *et al.* Heart disease and stroke statistics—2012 update: a report from the American Heart Association. *Circulation* 2012;125:e2–220.
- 2 Larsen MP, Eisenberg MS, Cummins RO, *et al.* Predicting survival from out-of-hospital cardiac arrest: a graphic model. *Ann Emerg Med* 1993;22:1652–8.
- 3 Link MS, Atkins DL, Passman RS, *et al.* Part 6: electrical therapies: automated external defibrillators, defibrillation, cardioversion, and pacing: 2010 American Heart Association Guidelines for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care. *Circulation* 2010;122:S706–19.
- 4 Caffrey SL, Willoughby PJ, Pepe PE, *et al.* Public use of automated external defibrillators. *N Engl J Med* 2002;347:1242–7.
- 5 Hallstrom AP, Ornato JP, Weisfeldt M, *et al.* Public-access defibrillation and survival after out-of-hospital cardiac arrest. *N Engl J Med* 2004;351:637–46.
- 6 Page RL, Joglar JA, Kowal RC, *et al.* Use of automated external defibrillators by a U.S. airline. *N Engl J Med* 2000;343:1210–16.
- 7 Valenzuela TD, Roe DJ, Nichol G, *et al.* Outcomes of rapid defibrillation by security officers after cardiac arrest in casinos. *N Engl J Med* 2000;343:1206–9.
- 8 Weisfeldt ML, Sitlani CM, Ornato JP, *et al.* Survival after application of automatic external defibrillators before arrival of the emergency medical system: evaluation in the resuscitation outcomes consortium population of 21 million. *J Am Coll Cardiol* 2010;55:1713–20.
- 9 Berdowski J, Blom MT, Bardai A, *et al.* Impact of onsite or dispatched automated external defibrillator use on survival after out-of-hospital cardiac arrest. *Circulation* 2012;124:2225–32.
- 10 Harmon KG, Asif IM, Klossner D, *et al.* Incidence of sudden cardiac death in national collegiate athletic association athletes. *Circulation* 2011;123:1594–600.
- 11 Maron BJ, Doerer JJ, Haas TS, *et al.* Sudden deaths in young competitive athletes: analysis of 1866 deaths in the United States, 1980–2006. *Circulation* 2009;119:1085–92.
- 12 Corrado D, Basso C, Rizzoli G, *et al.* Does sports activity enhance the risk of sudden death in adolescents and young adults? *J Am Coll Cardiol* 2003;42:1959–63.
- 13 Marijon E, Tafflet M, Celermajer DS, *et al.* Sports-related sudden death in the general population. *Circulation* 2011;124:672–81.
- 14 Albert CM, Mittleman MA, Chae CU, *et al.* Triggering of sudden death from cardiac causes by vigorous exertion. *N Engl J Med* 2000;343:1355–61.
- 15 Thompson PD, Franklin BA, Balady GJ, *et al.* Exercise and acute cardiovascular events placing the risks into perspective: a scientific statement from the American Heart Association Council on Nutrition, Physical Activity, and Metabolism and the Council on Clinical Cardiology. *Circulation* 2007;115:2358–68.
- 16 Drezner JA, Courson RW, Roberts WO, *et al.* Inter-association task force recommendations on emergency preparedness and management of sudden cardiac arrest in high school and college athletic programs: a consensus statement. *Heart Rhythm* 2007;4:549–65.
- 17 Drezner JA, Rao AL, Heistand J, *et al.* Effectiveness of emergency response planning for sudden cardiac arrest in United States high schools with automated external defibrillators. *Circulation* 2009;120:518–25.
- 18 Drezner JA. Preparing for sudden cardiac arrest—the essential role of automated external defibrillators in athletic medicine: a critical review. *Br J Sports Med* 2009;43:702–7.
- 19 The American Heart Association in Collaboration with the International Liaison Committee on Resuscitation. Guidelines 2000 for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care. Part 4: the automated external defibrillator: key link in the chain of survival. *Circulation* 2000;102:160–76.
- 20 Becker LB, Ostrander MP, Barrett J, *et al.* Outcome of CPR in a large metropolitan area—where are the survivors? *Ann Emerg Med* 1991;20:355–61.
- 21 Gallagher EJ, Lombardi G, Gennis P. Effectiveness of bystander cardiopulmonary resuscitation and survival following out-of-hospital cardiac arrest. *JAMA* 1995;274:1922–5.
- 22 Bobrow BJ, Clark LL, Ewy GA, *et al.* Minimally interrupted cardiac resuscitation by emergency medical services for out-of-hospital cardiac arrest. *JAMA* 2008;299:1158–65.
- 23 Toresdahl B, Harmon K, Drezner J. High school AED programs as a marker of emergency preparedness for sudden cardiac arrest. *J Athl Train* 2013;48:242–7.
- 24 Hazinski MF, Markenson D, Neish S, *et al.* Response to cardiac arrest and selected life-threatening medical emergencies: the medical emergency response plan for

- schools: a statement for healthcare providers, policymakers, school administrators, and community leaders. *Circulation* 2004;109:278–91.
- 25 Drezner J, Pluim B, Engebretsen L. Prevention of sudden cardiac death in athletes: new data and modern perspectives confront challenges in the 21st century. *Br J Sports Med* 2009;43:625–6.
- 26 Mahle WT, Sable CA, Matherne PG, *et al.* Key concepts in the evaluation of screening approaches for heart disease in children and adolescents: a science advisory from the American Heart Association. *Circulation* 2012;125:2796–801.
- 27 Kaltman JR, Thompson PD, Lantos J, *et al.* Screening for sudden cardiac death in the young: report from a national heart, lung, and blood institute working group. *Circulation* 2011;123:1911–18.
- 28 Campbell RM, Berger S, Drezner J. Sudden cardiac arrest in children and young athletes: the importance of a detailed personal and family history in the pre-participation evaluation. *Br J Sports Med* 2009;43:336–41.
- 29 Drezner JA, Rogers KJ. Sudden cardiac arrest in intercollegiate athletes: detailed analysis and outcomes of resuscitation in nine cases. *Heart Rhythm* 2006;3:755–9.
- 30 Drezner JA, Chun JS, Harmon KG, *et al.* Survival trends in the United States following exercise-related sudden cardiac arrest in the youth: 2000–2006. *Heart Rhythm* 2008;5:794–9.
- 31 Rothmier JD, Drezner JA, Harmon KG. Automated external defibrillators in Washington State high schools. *Br J Sports Med* 2007;41:301–5.
- 32 Lotfi K, White L, Rea T, *et al.* Cardiac arrest in schools. *Circulation* 2007;116:1374–9.
- 33 Drezner JA, Rogers KJ, Zimmer RR, *et al.* Use of automated external defibrillators at NCAA Division I Universities. *Med Sci Sports Exerc* 2005;37:1487–92.
- 34 Drezner JA, Rogers KJ, Horneff JG. Automated external defibrillator use at NCAA Division II and III universities. *Br J Sports Med* 2012;45:1174–8.